



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|--|
| (51) International Patent Classification ⁷ : A61K 45/06, A61P 9/00 | A1 | (11) International Publication Number: WO 00/38721 (43) International Publication Date: 6 July 2000 (06.07.00) |
| (21) International Application Number: PCT/US99/27942 (22) International Filing Date: 17 December 1999 (17.12.99) (30) Priority Data: 60/113,955 23 December 1998 (23.12.98) US 60/142,684 7 July 1999 (07.07.99) US (71) Applicant (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Department, P.O. Box 5110, Chicago, IL 60680-5110 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): SIKORSKI, James, A. [US/US]; 2313 East Royal Court, Des Peres, MO 63131 (US). GLENN, Kevin, C. [US/US]; 509 Princeton Gate Court, Chesterfield, MO 63017 (US). (74) Agents: WILLIAMS, Roger, A. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US). | | (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |
| (54) Title: COMBINATIONS OF CHOLESTERYL ESTER TRANSFER PROTEIN INHIBITORS AND NICOTINIC ACID DERIVATIVES FOR CARDIOVASCULAR INDICATIONS | | |
| (57) Abstract The present invention provides combinations of cardiovascular therapeutic compounds for the prophylaxis or treatment of cardiovascular disease including hypercholesterolemia, atherosclerosis, or hyperlipidemia. Combinations disclosed include a nicotinic acid derivative combined with a cholesteryl ester transfer protein (CETP) inhibitor. | | |

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

Combinations of Cholesteryl Ester Transfer Protein
Inhibitors and Nicotinic Acid Derivatives for
Cardiovascular Indications

5 This application claims priority of U.S. provisional application Ser. No. 60/142,684 filed Jul. 7, 1999 and of U.S. provisional application Ser. No. 60/113,955 filed Dec. 23, 1998.

10

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention relates to methods of treating cardiovascular diseases, and specifically relates to
15 combinations of compounds, compositions, and methods for their use in medicine, particularly in the prophylaxis and treatment of hyperlipidemic conditions such as are associated with atherosclerosis, hypercholesterolemia, and other coronary artery disease in mammals. More
20 particularly, the invention relates to cholesteryl ester transfer protein (CETP) activity inhibitors. The invention also relates to nicotinic acid derivatives.

Description of Related Art

25 It is well-settled that hyperlipidemic conditions associated with elevated concentrations of total cholesterol and low-density lipoprotein (LDL) cholesterol are major risk factors for coronary heart disease and particularly atherosclerosis. Numerous
30 studies have demonstrated that a low plasma concentration of high density lipoprotein (HDL) cholesterol is a powerful risk factor for the development of atherosclerosis (Barter and Rye, Atherosclerosis, 121, 1-12 (1996)). HDL is one of the

major classes of lipoproteins that function in the transport of lipids through the blood. The major lipids found associated with HDL include cholesterol, cholesteryl ester, triglycerides, phospholipids and
5 fatty acids. The other classes of lipoproteins found in the blood are low density lipoprotein (LDL), intermediate density lipoprotein (IDL), and very low density lipoprotein (VLDL). Since low levels of HDL cholesterol increase the risk of atherosclerosis,
10 methods for elevating plasma HDL cholesterol would be therapeutically beneficial for the treatment of atherosclerosis and other diseases associated with accumulation of lipid in the blood vessels. These diseases include, but are not limited to, coronary heart
15 disease, peripheral vascular disease, and stroke.

Atherosclerosis underlies most coronary artery disease (CAD), a major cause of morbidity and mortality in modern society. High LDL cholesterol (above about 180
mg/dl) and low HDL cholesterol (below 35 mg/dl) have been
20 shown to be important contributors to the development of atherosclerosis. Other diseases or risk factors, such as peripheral vascular disease, stroke, and hypercholesterolaemia are negatively affected by adverse HDL/LDL ratios.

25 Interfering with the recirculation of bile acids from the lumen of the intestinal tract is found to reduce the levels of serum cholesterol in a causal relationship. Epidemiological data has accumulated which indicates such reduction leads to an improvement in the disease state of
30 atherosclerosis. Stedronsky, in "Interaction of bile acids and cholesterol with nonsystemic agents having hypocholesterolemic properties," Biochimica et Biophysica Acta, 1210, 255-287 (1994) discusses the biochemistry,

physiology and known active agents surrounding bile acids and cholesterol.

Inhibition of cholesteryl ester transfer protein (CETP) has been shown to effectively modify plasma HDL/LDL ratios, and is expected to check the progress and/or formation of certain cardiovascular diseases. CETP is a plasma protein that facilitates the movement of cholesteryl esters and triglycerides between the various lipoproteins in the blood (Tall, J. Lipid Res., 34, 1255-74 (1993)). The movement of cholesteryl ester from HDL to LDL by CETP has the effect of lowering HDL cholesterol. It therefore follows that inhibition of CETP should lead to elevation of plasma HDL cholesterol and lowering of plasma LDL cholesterol, thereby providing a therapeutically beneficial plasma lipid profile. Evidence of this effect is described in McCarthy, Medicinal Res. Revs., 13, 139-59 (1993). Further evidence of this effect is described in Sitori, Pharmac. Ther., 67, 443-47 (1995)). This phenomenon was first demonstrated by Swenson et al., (J. Biol. Chem., 264, 14318 (1989)) with the use of a monoclonal antibody that specifically inhibits CETP. In rabbits, the antibody caused an elevation of the plasma HDL cholesterol and a decrease in LDL cholesterol. Son et al. (Biochim. Biophys. Acta, 795, 743-480 (1984)) describe proteins from human plasma that inhibit CETP. U.S. Patent 5,519,001, herein incorporated by reference, issued to Kushwaha et al., describes a 36 amino acid peptide derived from baboon apo C-1 that inhibits CETP activity. Cho et al. (Biochim. Biophys. Acta 1391, 133-144 (1998)) describe a peptide from hog plasma that inhibits human CETP. Bonin et al. (J. Peptide Res., 51, 216-225 (1998)) disclose a decapeptide inhibitor of

CETP. A depspeptide fungal metabolite is disclosed as a CETP inhibitor by Hedge et al. in *Bioorg. Med. Chem. Lett.*, 8, 1277-80 (1998).

There have been several reports of non-peptidic compounds that act as CETP inhibitors. Barrett et al. (*J. Am. Chem. Soc.*, 118, 7863-63 (1996)) describe cyclopropane-containing CETP inhibitors. Further cyclopropane-containing CETP inhibitors are described by Kuo et al. (*J. Am. Chem. Soc.*, 117, 10629-34 (1995)).

10 Pietzonka et al. (*Bioorg. Med. Chem. Lett.*, 6, 1951-54 (1996)) describe phosphonate-containing analogs of cholesteryl ester as CETP inhibitors. Coval et al. (*Bioorg. Med. Chem. Lett.*, 5, 605-610 (1995)) describe Wiedendiol-A and -B, and related sesquiterpene compounds

15 as CETP inhibitors. Lee et al. (*J. Antibiotics*, 49, 693-96 (1996)) describe CETP inhibitors derived from an insect fungus. Busch et al. (*Lipids*, 25, 216-220, (1990)) describe cholesteryl acetyl bromide as a CETP

inhibitor. Morton and Zilversmit (*J. Lipid Res.*, 35, 20 836-47 (1982)) describe that p-chloromercuriphenyl sulfonate, p-hydroxymercuribenzoate and ethyl mercurithiosalicylate inhibit CETP. Connolly et al. (*Biochem. Biophys. Res. Comm.*, 223, 42-47 (1996)) describe other cysteine modification reagents as CETP

25 inhibitors. Xia et al. describe 1,3,5-triazines as CETP inhibitors (*Bioorg. Med. Chem. Lett.*, 6, 919-22 (1996)). Bisgaier et al. (*Lipids*, 29, 811-8 (1994)) describe 4-phenyl-5-tridecyl-4H-1,2,4-triazole-thiol as a CETP inhibitor. Additional triazole CETP inhibitors

30 are described in U.S. Patent Application Serial No. 09/153,360, herein incorporated by reference. Sikorski et al. disclosed further novel CETP inhibitors in PCT Patent Application No. WO 9914204.

Substituted 2-mercaptoaniline amide compounds can be used as CETP inhibitors and such therapeutic compounds are described by H. Shinkai et al. in PCT Patent Application No. WO 98/35937.

- 5 Some substituted heteroalkylamine compounds are known as CETP inhibitors. In European Patent Application No. 796846, Schmidt et al. describe 2-aryl-substituted pyridines as cholesterol ester transfer protein inhibitors useful as cardiovascular agents. One
- 10 substituent at C₃ of the pyridine ring can be an hydroxyalkyl group. In European Patent Application No. 801060, Dow and Wright describe heterocyclic derivatives substituted with an aldehyde addition product of an alkylamine to afford 1-hydroxy-1-amines. These are
- 15 reported to be β 3-adrenergic receptor agonists useful for treating diabetes and other disorders. In Great Britain Patent Application No. 2305665, Fisher et al.
-
- ~~disclose 3-agonist-secondary-amino-alcohol-substituted~~
- pyridine derivatives useful for treating several
- 20 disorders including cholesterol levels and atherosclerotic diseases. In European Patent Application No. 818448 (herein incorporated by reference), Schmidt et al. describe tetrahydroquinoline derivatives as cholesterol ester transfer protein
- 25 inhibitors. European Patent Application No. 818197, Schmek et al. describe pyridines with fused heterocycles as cholesterol ester transfer protein inhibitors. Brandes et al. in German Patent Application No. 19627430 describe bicyclic condensed pyridine derivatives as
- 30 cholesterol ester transfer protein inhibitors. In PCT Patent Application No. WO 9839299, Muller-Gliemann et al. describe quinoline derivatives as cholesteryl ester transfer protein inhibitors.

Polycyclic compounds that are useful as CETP inhibitors are also disclosed by A. Oomura et al. in Japanese Patent No. 10287662. For example, therapeutic compounds having the structures C-1 and C-8 were
5 prepared by culturing *Penicillium* spp.

Cycloalkylpyridines useful as CETP inhibitors are disclosed by Schmidt et al. in European Patent No. EP 818448. For example, the therapeutic compound having the structure C-9 is disclosed as being particularly
10 effective as a CETP inhibitor.

Substituted tetrahydronaphthalene compounds useful as CETP inhibitors are described in PCT Patent Application No. WO 9914174. Specifically described in that disclosure as a useful CETP inhibitor is (8S)-3-
15 cyclopentyl-1-(4-fluorophenyl)-2-[(S)-fluoro(4-trifluoromethylphenyl)methyl]-8-hydroxy-6-spirocclobutyl-5,6,7,8-tetrahydronaphthalene.

Some 4-heteroaryl-tetrahydroquinolines useful as CETP inhibitors are described in PCT Patent Application
20 No. WO 9914215. For example, that disclosure describes 3-(4-trifluoromethylbenzoyl)-5,6,7,8-tetrahydroquinolin-5-one as a useful CETP inhibitor.

Nicotinic acid (niacin) is a B-complex vitamin reported as early as 1955 to act as a hypolipidemic agent
25 (R. Altschl, et al., Arch. Biochem. Biophys., 54, 558-9 (1955)). It is sometimes used to raise low HDL levels and lower VLDL and LDL levels. Useful commercial formulations of nicotinic acid include Niacor, Niaspan, Nicobid, Nicolar, Slo-Niacin. Nicotinic acid is contraindicated
30 for patients having hepatic dysfunction, active peptic ulcer, or arterial bleeding. Another compound in this class useful for cardiovascular indications is niceritrol (T. Kazumi et al., Curr. Ther. Res., 55, 546-51). J.

Sasaki et al. (Int. J. Clin. Pharm. Ther., 33 (7), 420-26 (1995)) describes a reduction in cholesteryl ester transfer activity by niceritrol monotherapy. Acipimox (5-methyl pyrazine-2-carboxylic acid 4-oxide, U.S. Patent No. 4,002,750) is structurally similar to nicotinic acid and has antihyperlipidemic activity.

Some combination therapies for the treatment of cardiovascular disease have been described in the literature. Combinations of IBAT inhibitors with HMG CoA reductase inhibitors useful for the treatment of cardiovascular disease are disclosed in U.S. Patent Application No. 09/037,308.

A combination therapy of fluvastatin and niceritrol is described by J. Sasaki et al. (Id.). Those researchers conclude that the combination of fluvastatin with niceritrol "at a dose of 750 mg/day dose does not appear to augment or attenuate beneficial effects of fluvastatin."

L. Cashin-Hemphill et al. (J. Am. Med. Assoc., 264 (23), 3013-17 (1990)) describe beneficial effects of a combination therapy of colestipol and niacin on coronary atherosclerosis. The described effects include nonprogression and regression in native coronary artery lesions.

A combination therapy of acipimox and simvastatin shows beneficial HDL effects in patients having high triglyceride levels (N. Hoogerbrugge et al., J. Internal Med., 241, 151-55 (1997)).

Sitostanol ester margarine and pravastatin combination therapy is described by H. Gylling et al. (J. Lipid Res., 37, 1776-85 (1996)). That therapy is reported to simultaneously inhibit cholesterol absorption and lower

LDL cholesterol significantly in non-insulin-dependent diabetic men.

Brown et al. (New Eng. J. Med., 323 (19), 1289-1339 (1990)) describe a combination therapy of lovastatin and
5 colestipol which reduces atherosclerotic lesion progression and increase lesion regression relative to lovastatin alone.

A combination therapy of an apoB secretion inhibitor with a CETP inhibitor was disclosed by Chang et al. in PCT
10 Patent Application No. WO 9823593.

Buch et al. (PCT Patent Application No. WO 9911263) describe a combination therapy comprising amlodipine and a statin compound for treating subjects suffering from angina pectoris, atherosclerosis, combined hypertension
15 and hyperlipidemia, and to treat symptoms of cardiac arrest. Buch et al. describe in PCT Patent Application No. WO 9911259 a combination therapy comprising amlodipine and atorvastatin.

Scott et al. (PCT Patent Application No. WO 9911260)
20 describe a combination therapy comprising atorvastatin and an antihypertensive agent.

Dettmar and Gibson (UK Patent Application No. GB 2329334 A) claim a therapeutic composition useful for reducing plasma low density lipoprotein and cholesterol
25 levels, wherein the composition comprises an HMG CoA reductase inhibitor and a bile complexing agent.

The above references show continuing need to find safe, effective agents for the prophylaxis or treatment of cardiovascular diseases.

30

Summary of the Invention

To address the continuing need to find safe and effective agents for the prophylaxis and treatment of

cardiovascular diseases, combination therapies of cardiovascular drugs are now reported.

Among its several embodiments, the present invention provides a combination therapy comprising the use of a
5 first amount of an CETP inhibitor and a second amount of another cardiovascular therapeutic useful in the prophylaxis or treatment of hyperlipidemia, atherosclerosis, or hypercholesterolemia, wherein said first and second amounts together comprise an anti-
10 hyperlipidemic condition effective amount, an anti-atherosclerotic condition effective amount, or an anti-hypercholesterolemic condition effective amount of the compounds. For example one of the many embodiments of the present invention is a combination therapy comprising
15 therapeutic dosages of an CETP inhibitor and a nicotinic acid derivative (including nicotinic acid or niacin).

A further embodiment of the instant invention comprises the use of any of the cardiovascular combination
therapies described herein for the prophylaxis or
20 treatment of hypercholesterolemia, atherosclerosis, or hyperlipidemia. Therefore, in one embodiment the present invention provides a method for the prophylaxis or treatment of a hyperlipidemic condition comprising administering to a patient in need thereof a combination
25 in unit dosage form wherein the combination comprises a first amount of a nicotinic acid derivative compound and a second amount of a CETP inhibiting compound wherein the first amount and the second amount together comprise an anti-hyperlipidemic condition effective amount of the
30 compounds.

In another embodiment, the present invention provides a method for the prophylaxis or treatment of an atherosclerotic condition comprising administering to a

patient in need thereof a combination in unit dosage form wherein the combination comprises a first amount of an nicotinic acid derivative compound and a second amount of a CETP inhibiting compound wherein the first amount and
5 the second amount together comprise an anti-atherosclerotic condition effective amount of the compounds.

In still another embodiment, the present invention provides method for the prophylaxis or treatment of
10 hypercholesterolemia comprising administering to a patient in need thereof a combination in unit dosage form wherein the combination comprises a first amount of an nicotinic acid derivative compound and a second amount of a CETP inhibiting compound wherein the first amount and the
15 second amount together comprise an anti-hypercholesterolemic condition effective amount of the compounds.

Further scope of the applicability of the present invention will become apparent from the detailed
20 description provided below. However, it should be understood that the following detailed description and examples, while indicating preferred embodiments of the invention, are given by way of illustration only since various changes and modifications within the spirit and
25 scope of the invention will become apparent to those skilled in the art from this detailed description.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

30 The following detailed description is provided to aid those skilled in the art in practicing the present invention. Even so, this detailed description should not be construed to unduly limit the present invention as

modifications and variations in the embodiments discussed herein can be made by those of ordinary skill in the art without departing from the spirit or scope of the present inventive discovery.

5 The contents of each of the references cited herein, including the contents of the references cited within these primary references, are herein incorporated by reference in their entirety.

10 a. Definitions

 The following definitions are provided in order to aid the reader in understanding the detailed description of the present invention: "

 "Nicotinic acid derivative" or "nicotinic acid
15 derivative compound" means a therapeutic compound comprising a pyridine-3-carboxylate structure or a pyrazine-2-carboxylate structure, including acid forms, salts, esters, zwitterions, and tautomers. Nicotinic acid
derivatives include, for example, nicotinic acid (niacin),
20 niceritrol, and acipimox.

 As used herein the term "CETP inhibitor" or "CETP inhibiting compound" means any therapeutic compound derived from chemical or biological sources which inhibits cholesteryl ester transfer protein activity.

25 "Combination therapy" means the administration of two or more therapeutic agents to treat a hyperlipidemic condition, for example atherosclerosis and hypercholesterolemia. Such administration encompasses co-administration of these therapeutic agents in a
30 substantially simultaneous manner, such as in a single capsule having a fixed ratio of active ingredients or in multiple, separate capsules for each inhibitor agent. In addition, such administration also encompasses use of each

type of therapeutic agent in a sequential manner. In either case, the treatment regimen will provide beneficial effects of the drug combination in treating the hyperlipidemic condition.

5 The phrase "therapeutically effective" is intended to qualify the combined amount of inhibitors in the combination therapy. This combined amount will achieve the goal of reducing or eliminating the hyperlipidemic condition.

10 "Therapeutic compound" means a compound useful in the prophylaxis or treatment of a hyperlipidemic condition, including atherosclerosis and hypercholesterolemia.

b. Combinations

15

The combinations of the present invention will have a number of uses. For example, through dosage adjustment and medical monitoring, the individual dosages of the therapeutic compounds used in the combinations of the

20 present invention will be lower than are typical for dosages of the therapeutic compounds when used in monotherapy. The dosage lowering will provide advantages including reduction of side effects of the individual therapeutic compounds when compared to the monotherapy.

25 In addition, fewer side effects of the combination therapy compared with the monotherapies will lead to greater patient compliance with therapy regimens.

Compounds useful in the present invention encompass a wide range of therapeutic compounds. Some individual CETP
30 inhibitor compounds useful in the present invention are separately described in the following individual patent applications, each of which is herein incorporated by reference.





- 5 R9. U.S. Patent Application Serial No. 60/101661.
R10. U.S. Patent Application Serial No. 60/101711.
R11. Patent Application Serial No. 60/101660.
R12. Patent Application Serial No. 60/101664.
R13. Patent Application Serial No. 60/101668.
R14. Patent Application Serial No. 60/101662.
R15. Patent Application Serial No. 60/101663.
R16. Patent Application Serial No. 60/101669.
10 R17. Patent Application Serial No. 60/101667.
R18. U.S. Patent Application Serial No. 09/401,916.
R19. U.S. Patent Application Serial No. 09/405,524.
R20. U.S. Patent Application Serial No. 09/404,638.
R21. U.S. Patent Application Serial No. 09/404,638.
15 R22. U.S. Patent Application Serial No. 09/400,915.
R23. U.S. Patent No. 5,932,587.
R24. U.S. Patent No. 5,925,645.




20 CETP inhibitor compounds of particular interest in
the present invention include those shown in Table 1, as
well as the diastereomers, enantiomers, racemates, salts,
and tautomers of the CETP inhibitors of Table 1.

Table 1.

25





| Compound Number | Structure |
|--------------------|-----------|
|--------------------|-----------|


| | |
|-----|--|
| C-1 |  |
| C-2 |  |
| C-3 |  |
| C-4 |  |

| | |
|-----|---|
| C-5 |  <chem>CCCCCCCCCCCCCN1C(=N)N(C1=S)c2ccccc2F</chem> |
| C-6 |  <chem>CCCCCCCCCCCCCN1C(=N)N(C1=S)c2ccccc2C</chem> |
| C-7 |  <chem>CCOc1ccc(Cl)cc1Oc2ccc(cc2)N(Cc3ccc(cc3)C(F)(F)F)C(C(F)(F)F)O</chem> |

| | |
|------|--|
| C-8 | |
| C-9 | |
| C-10 | |

| | |
|------|--|
| C-11 | <chem>CC(F)(O)CN(Cc1ccc(OCC(F)F)cc1)C2=CC=C(C=C2)C3=CC=C(C=C3)OCC(F)(F)F</chem> |
| C-12 | <chem>CC(F)(O)CN(Cc1ccc(OCC(F)F)cc1)C2=CC=C(C=C2)C3=CC=C(C=C3)OCC4=CC=C(Cl)C=C4</chem> |
| C-13 | <chem>CC(F)(O)CN(Cc1ccc(OCC(F)F)cc1)C2=CC=C(C=C2)C3=CC=C(C=C3)OCC4=CC=CC=C4</chem> |
| C-14 | <chem>CC(F)(O)CN(Cc1ccc(OCC(F)F)cc1)C2=CC=C(C=C2)C3=CC=C(C=C3)OCC4=CC=C(CC)C=C4</chem> |
| C-15 | |

| | |
|------|--|
| |  |
| C-16 |  |
| C-17 |  |
| C-18 |  |



Nicotinic acid derivatives useful in the combinations and methods of the present invention comprise a wide variety of structures and functionalities. Preferred nicotinic acid derivatives for the present invention are described in Table 2. The therapeutic compounds of Table 2 can be used in the present invention in a variety of forms, including acid form, salt form, racemates, enantiomers, zwitterions, and tautomers. The individual patent documents referenced in Table 2 are each herein incorporated by reference.

Table 2.

| Compound Number | Common Name | CAS Registry Number | Patent Document Reference |
|-----------------|----------------|---------------------|---------------------------|
| G-118 | Nicotinic Acid | 59-67-6 | |
| G-117 | Niceritrol | 5868-05-3 | GB 1022880 |

| | | | |
|-----|----------|------------|------------|
| G-3 | Acipimox | 51037-30-0 | GB 1351967 |
|-----|----------|------------|------------|

The compounds (for example, nicotinic acid derivatives or CETP inhibiting compounds) useful in the present invention can have no asymmetric carbon atoms, or, alternatively, the useful compounds can have one or more asymmetric carbon atoms. When the useful compounds have one or more asymmetric carbon atoms, they therefore include racemates and stereoisomers, such as diastereomers and enantiomers, in both pure form and in admixture. Such stereoisomers can be prepared using conventional techniques, for example by reacting enantiomeric starting materials, or by separating isomers of compounds of the present invention.

Isomers may include geometric isomers, for example *cis*-isomers or *trans*-isomers across a double bond. All such isomers are contemplated among the compounds useful in the present invention.

The compounds useful in the present invention also include tautomers.

The compounds useful in the present invention as discussed below include their salts, solvates and prodrugs.

25

Dosages, Formulations, and Routes of Administration

The compositions of the present invention can be administered for the prophylaxis and treatment of hyperlipidemic diseases or conditions by any means, preferably oral, that produce contact of these compounds with their site of action in the body, for example in the

ileum, the plasma, or the liver of a mammal, e.g., a human.

For the prophylaxis or treatment of the conditions referred to above, the compounds useful in the

5 compositions and methods of the present invention can be used as the compound *per se*. Pharmaceutically acceptable salts are particularly suitable for medical applications because of their greater aqueous solubility relative to the parent compound. Such salts must clearly have a

10 pharmaceutically acceptable anion or cation. Suitable pharmaceutically acceptable acid addition salts of the compounds of the present invention when possible include those derived from inorganic acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric, sulfonic,

15 and sulfuric acids, and organic acids such as acetic, benzenesulfonic, benzoic, citric, ethanesulfonic, fumaric, gluconic, glycolic, isothionic, lactic, lactobionic, maleic, malic, methanesulfonic, succinic, toluenesulfonic, tartaric, and trifluoroacetic acids. The chloride salt is

20 particularly preferred for medical purposes. Suitable pharmaceutically acceptable base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, and alkaline earth salts such as magnesium and calcium salts.

25 The anions useful in the present invention are, of course, also required to be pharmaceutically acceptable and are also selected from the above list.

The compounds useful in the present invention can be presented with an acceptable carrier in the form of a

30 pharmaceutical composition. The carrier must, of course, be acceptable in the sense of being compatible with the other ingredients of the composition and must not be deleterious to the recipient. The carrier can be a solid

or a liquid, or both, and is preferably formulated with the compound as a unit-dose composition, for example, a tablet, which can contain from 0.05% to 95% by weight of the active compound. Other pharmacologically active

5 substances can also be present, including other compounds of the present invention. The pharmaceutical compositions of the invention can be prepared by any of the well known techniques of pharmacy, consisting essentially of admixing the components.

10 Optionally, the combination of the present invention can comprise a composition comprising a nicotinic acid derivative compound and a CETP inhibiting compound. In such a composition, the nicotinic acid derivative compound and the CETP inhibiting compound can be present in a
15 single dosage form, for example a pill, a capsule, or a liquid which contains both of the compounds.

These compounds can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic

20 compounds or as a combination of therapeutic compounds.

The amount of compound which is required to achieve the desired biological effect will, of course, depend on a number of factors such as the specific compound chosen, the use for which it is intended, the mode of
25 administration, and the clinical condition of the recipient.

Generally a total daily dose of a nicotinic acid derivative can be in the range of from about 500 to about 10,000 mg/day, preferably about 1000 to about 8000 mg/day,
30 and more preferably still about 3000 to about 6000 mg/day in single or divided doses.

For a CETP inhibitor, a total daily dose of about 0.01 to about 100 mg/kg body weight/day, and preferably

between about 0.5 to about 20 mg/kg body weight/day, may generally be appropriate.

The daily doses described in the preceding paragraphs for the various therapeutic compounds can be administered
5 to the patient in a single dose, or in proportionate multiple subdoses. Subdoses can be administered 2 to 6 times per day. Doses can be in sustained release form effective to obtain desired results.

In the case of pharmaceutically acceptable salts, the
10 weights indicated above refer to the weight of the acid equivalent or the base equivalent of the therapeutic compound derived from the salt.

Oral delivery of the combinations of the present invention can include formulations, as are well known in
15 the art, to provide prolonged or sustained delivery of the drug to the gastrointestinal tract by any number of mechanisms. These include, but are not limited to, pH sensitive release from the dosage form based on the

changing pH of the small intestine, slow erosion of a
20 tablet or capsule, retention in the stomach based on the physical properties of the formulation, bioadhesion of the dosage form to the mucosal lining of the intestinal tract, or enzymatic release of the active drug from the dosage form. For some of the therapeutic compounds useful in the
25 present invention (e.g., a CETP inhibitor or a nicotinic acid derivative), the intended effect is to extend the time period over which the active drug molecule is delivered to the site of action by manipulation of the dosage form. Thus, enteric-coated and enteric-coated
30 controlled release formulations are within the scope of the present invention. Suitable enteric coatings include cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate and anionic

polymers of methacrylic acid and methacrylic acid methyl ester.

The combinations of the present invention can be delivered orally either in a solid, in a semi-solid, or in
5 a liquid form. When in a liquid or in a semi-solid form, the combinations of the present invention can, for example, be in the form of a liquid, syrup, or contained in a gel capsule (e.g., a gel cap). In one embodiment, when a CETP inhibitor is used in a combination of the
10 present invention, the CETP inhibitor can be provided in the form of a liquid, syrup, or contained in a gel capsule. In another embodiment, when a nicotinic acid derivative is used in a combination of the present invention, the nicotinic acid derivative can be provided
15 in the form of a liquid, syrup, or contained in a gel capsule.

For a CETP inhibitor the intravenously administered dose can, for example, be in the range of from about 0.003
mg/kg body weight to about 1.0 mg/kg body weight,
20 preferably from about 0.01 mg/kg body weight to about 0.75 mg/kg body weight, more preferably from about 0.1 mg/kg body weight to about 0.6 mg/kg body weight.

When administered intravenously, the dose for a nicotinic acid derivative can, for example, be in the
25 range of from about 150 mg/kg body weight to about 3000 mg/kg body weight, preferably from about 300 mg/kg body weight to about 2000 mg/kg body weight, more preferably from about 500 mg/kg body weight to about 1000 mg/kg body weight.

30 The dose of any of these therapeutic compounds can be conveniently administered as an infusion of from about 10 ng/kg body weight to about 100 ng/kg body weight per minute. Infusion fluids suitable for this purpose can

contain, for example, from about 0.1 ng to about 10 mg, preferably from about 1 ng to about 10 mg per milliliter. Unit doses can contain, for example, from about 1 mg to about 10 g of the compound of the present invention.

5 Thus, ampoules for injection can contain, for example, from about 1 mg to about 100 mg.

Pharmaceutical compositions according to the present invention include those suitable for oral, rectal, topical, buccal (e.g., sublingual), and parenteral (e.g.,
10 subcutaneous, intramuscular, intradermal, or intravenous) administration, although the most suitable route in any given case will depend on the nature and severity of the condition being treated and on the nature of the particular compound which is being used. In most cases,
15 the preferred route of administration is oral.

Pharmaceutical compositions suitable for oral administration can be presented in discrete units, such as capsules, cachets, lozenges, or tablets, each containing a
predetermined amount of at least one therapeutic compound
20 useful in the present invention; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water or water-in-oil emulsion. As indicated, such compositions can be prepared by any suitable method of pharmacy which includes the step of
25 bringing into association the active compound(s) and the carrier (which can constitute one or more accessory ingredients). In general, the compositions are prepared by uniformly and intimately admixing the active compound with a liquid or finely divided solid carrier, or both,
30 and then, if necessary, shaping the product. For example, a tablet can be prepared by compressing or molding a powder or granules of the compound, optionally with one or more accessory ingredients. Compressed tablets can be

prepared by compressing, in a suitable machine, the compound in a free-flowing form, such as a powder or granules optionally mixed with a binder, lubricant, inert diluent and/or surface active/dispersing agent(s). Molded
5 tablets can be made by molding, in a suitable machine, the powdered compound moistened with an inert liquid diluent.

Pharmaceutical compositions suitable for buccal (sub-lingual) administration include lozenges comprising a compound of the present invention in a flavored base,
10 usually sucrose, and acacia or tragacanth, and pastilles comprising the compound in an inert base such as gelatin and glycerin or sucrose and acacia.

Pharmaceutical compositions suitable for parenteral administration conveniently comprise sterile aqueous
15 preparations of a compound of the present invention. These preparations are preferably administered intravenously, although administration can also be effected by means of subcutaneous, intramuscular, or intradermal injection.

Such preparations can conveniently be prepared by admixing
20 the compound with water and rendering the resulting solution sterile and isotonic with the blood. Injectable compositions according to the invention will generally contain from 0.1 to 5% w/w of a compound disclosed herein.

Pharmaceutical compositions suitable for rectal
25 administration are preferably presented as unit-dose suppositories. These can be prepared by admixing a compound of the present invention with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.

30 Pharmaceutical compositions suitable for topical application to the skin preferably take the form of an ointment, cream, lotion, paste, gel, spray, aerosol, or oil. Carriers which can be used include petroleum jelly

(e.g., Vaseline), lanolin, polyethylene glycols, alcohols, and combinations of two or more thereof. The active compound is generally present at a concentration of from 0.1 to 50% w/w of the composition, for example, from 0.5 to 2%.

Transdermal administration is also possible. Pharmaceutical compositions suitable for transdermal administration can be presented as discrete patches adapted to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. Such patches suitably contain a compound of the present invention in an optionally buffered, aqueous solution, dissolved and/or dispersed in an adhesive, or dispersed in a polymer. A suitable concentration of the active compound is about 1% to 35%, preferably about 3% to 15%. As one particular possibility, the compound can be delivered from the patch by electrotransport or iontophoresis, for example, as described in Pharmaceutical Research, 3(6), 318 (1986).

In any case, the amount of active ingredient that can be combined with carrier materials to produce a single dosage form to be administered will vary depending upon the host treated and the particular mode of administration.

The solid dosage forms for oral administration including capsules, tablets, pills, powders, gel caps, and granules noted above comprise one or more compounds useful in the present invention admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate or solubilizing agents such as cyclodextrins. In the case of capsules, tablets,

powders, granules, gel caps, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration can
5 include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening,
10 flavoring, and perfuming agents.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or setting agents and suspending agents. The
15 sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable

vehicles and solvents that may be employed are water,
20 Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids
25 such as oleic acid find use in the preparation of injectables.

Pharmaceutically acceptable carriers encompass all the foregoing and the like.

In combination therapy, administration of two or more
30 of the therapeutic agents useful in the present invention may take place sequentially in separate formulations, or may be accomplished by simultaneous administration in a single formulation or separate formulations.

Administration may be accomplished by oral route, or by intravenous, intramuscular, or subcutaneous injections. The formulation may be in the form of a bolus, or in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more pharmaceutically-acceptable carriers or diluents, or a binder such as gelatin or hydroxypropylmethyl cellulose, together with one or more of a lubricant, preservative, surface active or dispersing agent.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension, or liquid. Capsules, tablets, etc., can be prepared by conventional methods well known in the art. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient or ingredients. Examples of dosage units are tablets or capsules. These may with advantage contain one or more therapeutic compound in an amount described above. For example, in the case of a CETP inhibitor, the dose range may be from about 0.01 mg to about 500 mg or any other dose, dependent upon the specific inhibitor, as is known in the art. In the case of a nicotinic acid derivative, the dose range may be from about 0.01 mg to about 500 mg or any other dose, dependent upon the specific inhibitor, as is known in the art.

The active ingredients may also be administered by injection as a composition wherein, for example, saline, dextrose, or water may be used as a suitable carrier. A suitable daily dose of each active therapeutic compound is one that achieves the same blood serum level as produced by oral administration as described above.

The therapeutic compounds may further be administered by any combination of oral/oral, oral/parenteral, or parenteral/parenteral route.

Pharmaceutical compositions for use in the treatment
5 methods of the present invention may be administered in oral form or by intravenous administration. Oral administration of the combination therapy is preferred. Dosing for oral administration may be with a regimen calling for single daily dose, or for a single dose every
10 other day, or for multiple, spaced doses throughout the day. The therapeutic compounds which make up the combination therapy may be administered simultaneously, either in a combined dosage form or in separate dosage forms intended for substantially simultaneous oral
15 administration. The therapeutic compounds which make up the combination therapy may also be administered sequentially, with either therapeutic compound being administered by a regimen calling for two-step ingestion.

Thus, a regimen may call for sequential administration of
20 the therapeutic compounds with spaced-apart ingestion of the separate, active agents. The time period between the multiple ingestion steps may range from a few minutes to several hours, depending upon the properties of each therapeutic compound such as potency, solubility,
25 bioavailability, plasma half-life and kinetic profile of the therapeutic compound, as well as depending upon the effect of food ingestion and the age and condition of the patient. Circadian variation of the target molecule concentration may also determine the optimal dose
30 interval. The therapeutic compounds of the combined therapy whether administered simultaneously, substantially simultaneously, or sequentially, may involve a regimen calling for administration of one therapeutic compound by

oral route and another therapeutic compound by intravenous route. Whether the therapeutic compounds of the combined therapy are administered by oral or intravenous route, separately or together, each such therapeutic compound
5 will be contained in a suitable pharmaceutical formulation of pharmaceutically-acceptable excipients, diluents or other formulations components. Examples of suitable pharmaceutically-acceptable formulations containing the therapeutic compounds for oral administration are given
10 above.

Treatment Regimen

The dosage regimen to prevent, give relief from, or ameliorate a disease condition having hyperlipidemia as an
15 element of the disease, e.g., atherosclerosis, or to protect against or treat further high cholesterol plasma or blood levels with the compounds and/or compositions of the present invention is selected in accordance with a
variety of factors. These include the type, age, weight,
20 sex, diet, and medical condition of the patient, the severity of the disease, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetics and toxicology profiles of the particular compound employed, whether a drug delivery
25 system is utilized, and whether the compound is administered as part of a drug combination. Thus, the dosage regimen actually employed may vary widely and therefore deviate from the preferred dosage regimen set forth above.

30 Initial treatment of a patient suffering from a hyperlipidemic condition can begin with the dosages indicated above. Treatment should generally be continued as necessary over a period of several weeks to several

months or years until the hyperlipidemic disease condition has been controlled or eliminated. Patients undergoing treatment with the compounds or compositions disclosed herein can be routinely monitored by, for example,

5 measuring serum LDL and total cholesterol levels by any of the methods well known in the art, to determine the effectiveness of the combination therapy. Continuous analysis of such data permits modification of the treatment regimen during therapy so that optimal effective

10 amounts of each type of therapeutic compound are administered at any point in time, and so that the duration of treatment can be determined as well. In this way, the treatment regimen/dosing schedule can be rationally modified over the course of therapy so that the

15 lowest amount of the therapeutic compounds which together exhibit satisfactory effectiveness is administered, and so that administration is continued only so long as is necessary to successfully treat the hyperlipidemic

condition.

20 A potential advantage of the combination therapy disclosed herein may be reduction of the amount of any individual therapeutic compound, or all therapeutic compounds, effective in treating hyperlipidemic conditions such as atherosclerosis and hypercholesterolemia.

25 One of the several embodiments of the present invention comprises a combination therapy comprising the use of a first amount of an CETP inhibitor and a second amount of another cardiovascular therapeutic useful in the prophylaxis or treatment of hyperlipidemia,

30 atherosclerosis, or hypercholesterolemia wherein said first and second amounts together comprise an anti-hyperlipidemic condition effective amount, an anti-atherosclerotic condition effective amount, or an anti-

hypercholesterolemic condition effective amount of said compounds. For example one of the many embodiments of the present invention is a combination therapy comprising therapeutic dosages of an CETP inhibitor and a nicotinic
5 acid derivative.

The embodiments of the present invention can comprise a combination therapy using two or more of the therapeutic compounds described or incorporated herein. The combination therapy can comprise two or more therapeutic
10 compounds from different classes of chemistry, e.g., a nicotinic acid derivative can be therapeutically combined with a CETP inhibitor. Therapeutic combinations can comprise more than two therapeutic compounds. For example, the therapy can comprise the use of a nicotinic
15 acid derivative, a CETP inhibitor, and a HMG CoA reductase inhibitor. Alternatively, two or more therapeutic compounds from the same class of chemistry can comprise the therapy, e.g. a combination therapy comprising two or
more nicotinic acid derivatives or two or more CETP
20 inhibitors.

A further embodiment of the instant invention comprises the use of any of the cardiovascular combination therapies described herein for the prophylaxis or treatment of hypercholesterolemia, atherosclerosis, or
25 hyperlipidemia.

The following non-limiting examples serve to illustrate various aspects of the present invention.

c. Examples

30

Table 3 illustrates examples of some combinations of the present invention wherein the combination comprises a first amount of a CETP inhibitor and a second amount of a nicotinic acid derivative, wherein said first and second

amounts together comprise an anti-hyperlipidemic condition effective amount or an anti-atherosclerotic condition effective amount of said compounds.

5

Table 3.

| Example Number | Component 1 | Component 2 |
|----------------|-------------|------------------------------------|
| 1 | C-1 | nicotinic acid (niacin) |
| 2 | C-2 | nicotinic acid (niacin) |
| 3 | C-3 | nicotinic acid (niacin) |
| 4 | C-4 | nicotinic acid (niacin) |
| 5 | C-5 | nicotinic acid (niacin) |
| 6 | C-6 | nicotinic acid (niacin) |
| 7 | C-7 | nicotinic acid (niacin) |
| 8 | C-8 | nicotinic acid (niacin) |
| 9 | C-9 | nicotinic acid (niacin) |
| 10 | C-10 | nicotinic acid (niacin) |
| 11 | C-11 | nicotinic acid (niacin) |
| 12 | C-12 | nicotinic acid (niacin) |
| 13 | C-13 | nicotinic acid (niacin) |
| 14 | C-14 | nicotinic acid (niacin) |
| 15 | C-15 | nicotinic acid (niacin) |
| 16 | C-16 | nicotinic acid (niacin) |
| 17 | C-17 | nicotinic acid (niacin) |
| 18 | C-18 | nicotinic acid (niacin) |
| 19 | C-19 | nicotinic acid (niacin) |
| 20 | C-20 | nicotinic acid (niacin) |
| 21 | C-1 | niceritrol |
| 22 | C-2 | niceritrol |
| 23 | C-3 | niceritrol |
| 24 | C-4 | niceritrol |
| 25 | C-5 | niceritrol |
| 26 | C-6 | niceritrol |
| 27 | C-7 | niceritrol |
| 28 | C-8 | niceritrol |
| 29 | C-9 | niceritrol |
| 30 | C-10 | niceritrol |
| 31 | C-11 | niceritrol |
| 32 | C-12 | niceritrol |

| | | |
|----|------|------------|
| 33 | C-13 | niceritrol |
| 34 | C-14 | niceritrol |
| 35 | C-15 | niceritrol |
| 36 | C-16 | niceritrol |
| 37 | C-17 | niceritrol |
| 38 | C-18 | niceritrol |
| 39 | C-19 | niceritrol |
| 40 | C-20 | niceritrol |
| 41 | C-1 | acipimox |
| 42 | C-2 | acipimox |
| 43 | C-3 | acipimox |
| 44 | C-4 | acipimox |
| 45 | C-5 | acipimox |
| 46 | C-6 | acipimox |
| 47 | C-7 | acipimox |
| 48 | C-8 | acipimox |
| 49 | C-9 | acipimox |
| 50 | C-10 | acipimox |
| 51 | C-11 | acipimox |
| 52 | C-12 | acipimox |
| 53 | C-13 | acipimox |
| 54 | C-14 | acipimox |
| 55 | C-15 | acipimox |
| 56 | C-16 | acipimox |
| 57 | C-17 | acipimox |
| 58 | C-18 | acipimox |
| 59 | C-19 | acipimox |
| 60 | C-20 | acipimox |

BIOLOGICAL ASSAYS

The utility of the combinations of the present
5 invention can be shown by the following assays. These
assays are performed in vitro and in animal models
essentially using procedures recognized to show the
utility of the present invention.

In Vitro Assay of compounds that inhibit IBAT-mediated uptake of [14 C]-Taurocholate (TC) in H14 Cells

Baby hamster kidney cells (BHK) transfected with the cDNA of human IBAT (H14 cells) are seeded at 60,000
5 cells/well in 96 well Top-Count tissue culture plates for assays run within in 24 hours of seeding, 30,000 cells/well for assays run within 48 hours, and 10,000 cells/well for assays run within 72 hours.

On the day of assay, the cell monolayer is gently
10 washed once with 100 μ l assay buffer (Dulbecco's Modified Eagle's medium with 4.5 g/L glucose + 0.2% (w/v) fatty acid free bovine serum albumin- (FAF)BSA). To each well 50 μ l of a two-fold concentrate of test compound in assay buffer is added along with 50 μ l of 6 μ M [14 C]-
15 taurocholate in assay buffer (final concentration of 3 μ M [14 C]-taurocholate). The cell culture plates are incubated 2 hours at 37°C prior to gently washing each well twice
with 100 μ l 4°C Dulbecco's phosphate-buffered saline (PBS) containing 0.2% (w/v) (FAF)BSA. The wells are then gently
20 washed once with 100 μ l 4°C PBS without (FAF)BSA. To each 200 μ l of liquid scintillation counting fluid is added, the plates are heat sealed and shaken for 30 minutes at room temperature prior to measuring the amount of radioactivity in each well on a Packard Top-Count
25 instrument.

In Vitro Assay of compounds that inhibit uptake of [14 C]-Alanine

The alanine uptake assay is performed in an identical
30 fashion to the taurocholate assay, with the exception that labeled alanine is substituted for the labeled taurocholate.

In Vivo Assay of compounds that inhibit Rat Ileal uptake
of [¹⁴C]-Taurocholate into Bile

(See "Metabolism of 3 α ,7 β -dihydroxy-7 α -methyl-5 β -
5 cholanoic acid and 3 α ,7 β -dihydroxy-7 α -methyl-5 β -cholanoic
acid in hamsters" in Biochimica et Biophysica Acta, 833,
196-202 (1985) by Une et al., herein incorporated by
reference.)

Male wistar rats (200-300 g) are to be anesthetized
10 with inactin @100 mg/kg. Bile ducts are to be cannulated
with a 10" length of PE10 tubing. The small intestine is
to be exposed and laid out on a gauze pad. A canulae
(1/8" luer lock, tapered female adapter) is inserted at 12
cm from the junction of the small intestine and the cecum.
15 A slit is cut at 4 cm from this same junction (utilizing a
8 cm length of ileum). Twenty milliliters of warm
Dulbecco's phosphate buffered saline, pH 6.5 (PBS) is to
be used to flush out the intestine segment. The distal
opening is cannulated with a 20 cm length of silicone
20 tubing (0.02" I.D. x 0.037" O.D.). The proximal cannulae
is hooked up to a peristaltic pump and the intestine is
washed for 20 min with warm PBS at 0.25 ml/min.
Temperature of the gut segment is to be monitored
continuously. At the start of the experiment, 2.0 ml of
25 control sample ([¹⁴C]-taurocholate @ 0.05 mCi/mL with 5 mM
non-radiolabeled taurocholate) is to be loaded into the
gut segment with a 3 ml syringe and bile sample collection
is begun. Control sample is infused at a rate of 0.25
ml/min for 21 min. Bile samples fractions are to be
30 collected every 3 minute for the first 27 minutes of the
procedure. After the 21 min of sample infusion, the ileal
loop is to be washed out with 20 ml of warm PBS (using a
30 ml syringe), and then the loop is to be washed out for

21 min with warm PBS at 0.25 ml/min. A second perfusion is initiated as described above but this with test compound being administered as well (21 min administration followed by 21 min of wash out) and bile sampled every 3 min for the first 27 min. If necessary, a third perfusion will be performed as above that typically contains the control sample.

Measurement of Hepatic Cholesterol Concentration (HEPATIC CHOL)

Liver tissue is to be weighed and homogenized in chloroform:methanol (2:1). After homogenization and centrifugation the supernatant is separated and dried under nitrogen. The residue is to be dissolved in isopropanol and the cholesterol content will be measured enzymatically, using a combination of cholesterol oxidase and peroxidase, as described by Allain, C. A. et al., Clin. Chem., 20, 470 (1974) (herein incorporated by reference).

20

Measurement of Hepatic HMG CoA-Reductase Activity (HMG COA)

Hepatic microsomes are to be prepared by homogenizing liver samples in a phosphate/sucrose buffer, followed by centrifugal separation. The final pelleted material is resuspended in buffer and an aliquot will be assayed for HMG CoA reductase activity by incubating for 60 minutes at 37° C in the presence of ¹⁴C-HMG-CoA (Dupont-NEN). The reaction is stopped by adding 6N HCl followed by centrifugation. An aliquot of the supernatant is separated, by thin-layer chromatography, and the spot corresponding to the enzyme product is scraped off the plate, extracted and radioactivity is determined by

scintillation counting. (Reference: Akerlund, J. and Bjorkhem, I. (1990) *J. Lipid Res.* 31, 2159).

Determination of Serum Cholesterol (SER.CHOL, HDL-CHOL,

5 TGI and VLDL + LDL)

Total serum cholesterol (SER.CHOL) are to be measured enzymatically using a commercial kit from Wako Fine Chemicals (Richmond, VA); Cholesterol C11, Catalog No. 276-64909. HDL cholesterol (HDL-CHOL) will be assayed
10 using this same kit after precipitation of VLDL and LDL with Sigma Chemical Co. HDL Cholesterol reagent, Catalog No. 352-3 (dextran sulfate method). Total serum triglycerides (blanked) (TGI) will be assayed enzymatically with Sigma Chemical Co. GPO-Trinder, Catalog
15 No. 337-B. VLDL and LDL (VLDL + LDL) cholesterol concentrations will be calculated as the difference between total and HDL cholesterol.

Measurement of Hepatic Cholesterol 7- α -Hydroxylase

20 Activity (7 α -OHase)

Hepatic microsomes are to be prepared by homogenizing liver samples in a phosphate/sucrose buffer, followed by centrifugal separation. The final pelleted material is resuspended in buffer and an aliquot will be assayed for
25 cholesterol 7- α -hydroxylase activity by incubating for 5 minutes at 37° C in the presence of NADPH. Following extraction into petroleum ether, the organic solvent is evaporated and the residue is dissolved in acetonitrile/methanol. The enzymatic product will be separated by
30 injecting an aliquot of the extract onto a C₁₈ reversed phase HPLC column and quantitating the eluted material using UV detection at 240nm. (Reference: Horton, J. D., et al. (1994) *J. Clin. Invest.* 93, 2084).

Rat Gavage Assay

Male Wister rats (275-300g) are to be administered IBAT inhibitors using an oral gavage procedure. Drug or
5 vehicle (0.2% TWEEN 80 in water) is administered once a day (9:00-10:00 a.m.) for 4 days at varying dosages in a final volume of 2 mL per kilogram of body weight. (TWEEN 80 is a 20 molar polyethyleneoxide sorbitan monooleate surfactant manufactured by ICI Specialty Chemicals,
10 Wilmington, Delaware, U.S.A.) Total fecal samples are collected during the final 48 hours of the treatment period and analyzed for bile acid content using an enzymatic assay as described below. Compound efficacy will be determined by comparison of the increase in fecal
15 bile acid (FBA) concentration in treated rats to the mean FBA concentration of rats in the vehicle group.

Measurement of Rat Fecal Bile Acid Concentration (FBA)

Total fecal output from individually housed rats is
20 to be collected for 24 or 48 hours, dried under a stream of nitrogen, pulverized, mixed, and weighed. Approximately 0.1 gram is weighed out and extracted into an organic solvent (butanol/water). Following separation and drying, the residue is dissolved in methanol and the amount of
25 bile acid present will be measured enzymatically using the 3 α -hydroxysteroid steroid dehydrogenase reaction with bile acids to reduce NAD. (see Mashige, F. et al. Clin. Chem., 27, 1352 (1981), herein incorporated by reference).

30 [3H]Taurocholate Uptake in Rabbit Brush Border Membrane Vesicles (BBMV)

Rabbit Ileal brush border membranes are to be prepared from frozen ileal mucosa by the calcium

precipitation method describe by Malathi et al.

(Biochimica Biophysica Acta, 554, 259 (1979), herein incorporated by reference). The method for measuring taurocholate is essentially as described by Kramer et al.

- 5 (Biochimica Biophysica Acta, 1111, 93 (1992), herein incorporated by reference) except the assay volume will be 200 μ l instead of 100 μ l. Briefly, at room temperature a 190 μ l solution containing 2 μ M [3 H]-taurocholate(0.75 μ Ci), 20 mM tris, 100 mM NaCl, 100 mM mannitol pH 7.4 is
- 10 incubated for 5 sec with 10 μ l of brush border membrane vesicles (60-120 μ g protein). The incubation is initiated by the addition of the BBMV while vortexing and the reaction is to be stopped by the addition of 5 ml of ice cold buffer (20 mM Hepes-tris, 150 mM KCl) followed
- 15 immediately by filtration through a nylon filter (0.2 μ m pore) and an additional 5 ml wash with stop buffer.

Acyl-CoA; Cholesterol Acyl Transferase (ACAT)

- 20 Hamster liver and rat intestinal microsomes are to be prepared from tissue as described previously (J. Biol. Chem., 255, 9098 (1980), herein incorporated by reference) and used as a source of ACAT enzyme. The assay will consist of a 2.0 ml incubation containing 24 μ M Oleoyl-CoA
- 25 (0.05 μ Ci) in a 50 mM sodium phosphate, 2 mM DTT pH 7.4 buffer containing 0.25 % BSA and 200 μ g of microsomal protein. The assay will be initiated by the addition of oleoyl-CoA. The reaction proceeds for 5 min at 37° C and will be terminated by the addition of 8.0 ml of
- 30 chloroform/ methanol (2:1). To the extraction is added 125 μ g of cholesterol oleate in chloroform methanol to act as a carrier and the organic and aqueous phases of the extraction are separated by centrifugation after thorough

vortexing. The chloroform phase is to be taken to dryness and then spotted on a silica gel 60 TLC plate and developed in hexane/ethyl ether (9:1). The amount of cholesterol ester formed will be determined by measuring the amount of radioactivity incorporated into the cholesterol oleate spot on the TLC plate with a Packard Instainmager.

Dog Model for Evaluating Lipid Lowering Drugs

10 Male beagle dogs, obtained from a vendor such as Marshall farms and weighing 6-12 kg are fed once a day for two hours and given water ad libitum. Dogs may be randomly assigned to a dosing groups consisting of 6 to 12 dogs each, such as: vehicle, i.g.; 1mg/kg, i.g.; 2mg/kg, i.g.;
15 4 mg/kg, i.g.; 2 mg/kg, p.o. (powder in capsule). Intra-gastric dosing of a therapeutic material dissolved in aqueous solution (for example, 0.2% Tween 80 solution [polyoxyethylene mono-oleate, Sigma Chemical Co., St.
Louis, MO]) may be done using a gavage tube. Prior to
20 initiating dosing, blood samples may be drawn from the cephalic vein in the morning before feeding in order to evaluate serum cholesterol (total and HDL) and triglycerides. For several consecutive days animals are dosed in the morning, prior to feeding. Animals are to be
25 allowed 2 hours to eat before any remaining food is removed. Feces are to be collected over a 2 day period at the end of the study and may be analyzed for bile acid or lipid content. Blood samples are also to be taken, at the end of the treatment period, for comparison with pre-study
30 serum lipid levels. Statistical significance will be determined using the standard student's T-test with $p < .05$.

Dog Serum Lipid Measurement

Blood is to be collected from the cephalic vein of fasted dogs in serum separator tubes (Vacutainer SST, 5 Becton Dickinson and Co., Franklin Lakes, NJ). The blood is centrifuged at 2000 rpm for 20 minutes and the serum decanted.

Total cholesterol may be measured in a 96 well format using a Wako enzymatic diagnostic kit (Cholesterol CII) 10 (Wako Chemicals, Richmond, VA), utilizing the cholesterol oxidase reaction to produce hydrogen peroxide which is measured colorimetrically. A standard curve from 0.5 to 10 µg cholesterol is to be prepared in the first 2 columns of the plate. The serum samples (20-40 µl, depending on 15 the expected lipid concentration) or known serum control samples are added to separate wells in duplicate. Water is added to bring the volume to 100 µl in each well. A 100 µl aliquot of color reagent is added to each well and the plates will be read at 500 nm after a 15 minute 20 incubation at 37 degrees centigrade.

HDL cholesterol may be assayed using Sigma kit No. 352-3 (Sigma Chemical Co., St. Louis, MO) which utilizes dextran sulfate and Mg ions to selectively precipitate LDL and VLDL. A volume of 150 µl of each serum sample is to 25 be added to individual microfuge tubes, followed by 15 µl of HDL cholesterol reagent (Sigma 352-3). Samples are to be mixed and centrifuged at 5000 rpm for 5 minutes. A 50 µl aliquot of the supernatant is to be then mixed with 200 µl of saline and assayed using the same procedure as for 30 total cholesterol measurement.

Triglycerides are to be measured using Sigma kit No. 337 in a 96 well plate format. This procedure will

measure glycerol, following its release by reaction of triglycerides with lipoprotein lipase. Standard solutions of glycerol (Sigma 339-11) ranging from 1 to 24 μg are to be used to generate the standard curve. Serum samples
5 (20-40 μl , depending on the expected lipid concentration) are added to wells in duplicate. Water is added to bring the volume to 100 μl in each well and 100 μl of color reagent was also added to each well. After mixing and a 15 minute incubation, the plates will be read at 540 nm
10 and the triglyceride values calculated from the standard curve. A replicate plate is also to be run using a blank enzyme reagent to correct for any endogenous glycerol in the serum samples.

15 Dog Fecal Bile Acid Measurement

Fecal samples may be collected to determine the fecal bile acid (FBA) concentration for each animal. Fecal
collections may be made during the final 48 hours of the study, for two consecutive 24 hour periods between 9:00 am
20 and 10:00 am each day, prior to dosing and feeding. The separate two day collections from each animal are to be weighed, combined and homogenized with distilled water in a processor (Cuisinart) to generate a homogeneous slurry. About 1.4 g of the homogenate is to be extracted in a
25 final concentration of 50% tertiary butanol/distilled water (2:0.6) for 45 minutes in a 37°C water bath and centrifuged for 13 minutes at 2000 x g. The concentration of bile acids (mmoles/day) may be determined using a 96-well enzymatic assay system (1,2). A 20 μl aliquot of the
30 fecal extract is to be added to two sets each of triplicate wells in a 96-well assay plate. A standardized sodium taurocholate solution and a standardized fecal extract solution (previously made from pooled samples and

characterized for its bile acid concentration) will also analyzed for assay quality control. Twenty-microliter aliquots of sodium taurocholate, serially diluted to generate a standard curve are similarly to be added to two
5 sets of triplicate wells. A 230 μ l reaction mixture containing 1M hydrazine hydrate, 0.1 M pyrophosphate and 0.46 mg/ml NAD is to be added to each well. A 50 μ l aliquot of 3 α -hydroxysteroid dehydrogenase enzyme (HSD; 0.8 units/ml) or assay buffer (0.1 M sodium pyrophosphate)
10 are then added to one of the two sets of triplicates. All reagents may be obtained from Sigma Chemical Co., St. Louis, MO. Following 60 minutes of incubation at room temperature, the optical density at 340nm will be measured and the mean of each set of triplicate samples will be
15 calculated. The difference in optical density \pm HSD enzyme is to be used to determine the bile acid concentration (mM) of each sample based on the sodium taurocholate standard curve. The bile acid concentration of the extract, the weight of the fecal homogenate (grams)
20 and the body weight of the animal are to be used to calculate the corresponding FBA concentration in mmol/kg/day for each animal. The mean FBA concentration (mmol/kg/day) of the vehicle group is to be subtracted from the FBA concentration of each treatment group to
25 determine the increase (delta value) in FBA concentration as a result of the treatment.

CETP ACTIVITY ASSAY IN HUMAN PLASMA (Tritiated
cholesteryl ester)

30

Blood is to be obtained from healthy volunteers. Blood is collected in tubes containing EDTA (EDTA plasma pool). The EDTA human plasma pool previously stored at -

20°C, is to be thawed at room temperature, and centrifuged for 5 minutes to remove any particulate matter. Tritiated HDL, radiolabeled in the cholesteryl ester moiety ($[^3\text{H}]$ CE-HDL) as described by Morton and Zilversmit (J. Biol. Chem., 256, 11992-95 (1981)), is to be added to the plasma to a final concentration of (25 $\mu\text{g/ml}$ cholesterol). Inhibitor compounds are to be added to the plasma as follows: Equal volumes of the plasma containing the $[^3\text{H}]$ CE-HDL (396 μl) are added by pipette into micro tubes (Titertube[®], Bio-Rad laboratories, Hercules, CA). Compounds, usually dissolved as 20-50 mM stock solutions in DMSO, are to be serially diluted in DMSO (or an alternative solvent in some cases, such as dimethylformamide or ethanol). Four μl of each of the serial dilutions of inhibitor compounds or DMSO alone are then added to each of the plasma tubes. The tubes are immediately mixed. Triplicate aliquots (100 μl) from each plasma tube are then transferred to wells of 96-well round-bottomed polystyrene microtiter plates (Corning, Corning, NY). Plates are sealed with plastic film and incubated at 37°C for 4 hours. Test wells are to contain plasma with dilutions of inhibitor compounds. Control wells are to contain plasma with DMSO alone. Blank wells are to contain plasma with DMSO alone that are left in the micro tubes at 4°C for the 4 hour incubation and are added to the microtiter wells at the end of the incubation period. VLDL and LDL are precipitated by the addition of 10 μl of precipitating reagent (1% (w/v) dextran sulfate (Dextralip50)/0.5 M magnesium chloride, pH 7.4) to all wells. The wells are mixed on a plate mixer and then incubated at ambient temperature for 10 min. The plates are then centrifuged at 1000 x g for 30 min at 10°C. The

supernatants (50 μ l) from each well are then transferred to PicoplateTM 96 plate wells (Packard, Meriden, CT) containing 250:1 MicroscintTM-40 (Packard, Meriden, CT). The plates are heat-sealed (TopSealTM-P, Packard, Meriden, CT) according to the manufacturer's directions and mixed for 30 min. Radioactivity will be measured on a microplate scintillation counter (TopCount, Packard, Meriden, CT). IC₅₀ values will be determined as the concentration of inhibitor compound inhibiting transfer of [3H]CE from the supernatant [3H]CE-HDL to the precipitated VLDL and LDL by 50% compared to the transfer obtained in the control wells. The maximum percentage transfer (in the control wells) will be determined using the following equation:

15

$$\% \text{ Transfer} = \frac{[\text{dpm}_{\text{blank}} - \text{dpm}_{\text{control}}] \times 100}{\text{dpm}_{\text{blank}}}$$

The percentage of control transfer determined in the wells containing inhibitor compounds is determined as follows:

20

$$\% \text{ Control} = \frac{[\text{dpm}_{\text{blank}} - \text{dpm}_{\text{test}}] \times 100}{\text{dpm}_{\text{blank}} - \text{dpm}_{\text{control}}}$$

IC₅₀ values will be calculated from plots of % control versus concentration of inhibitor compound.

25

CETP Activity In Vitro

The ability of compounds to inhibit CETP activity are assessed using an in vitro assay that measures the rate of transfer of radiolabeled cholesteryl ester ([3H]CE) from

HDL donor particles to LDL acceptor particles. Details of the assay are provided by Glenn et al. (Glenn and Melton, "Quantification of Cholesteryl Ester Transfer Protein (CETP): A) CETP Activity and B) Immunochemical Assay of CETP Protein," Meth. Enzymol., 263, 339-351 (1996)). CETP can be obtained from the serum-free conditioned medium of CHO cells transfected with a cDNA for CETP (Wang, S. et al. J. Biol. Chem. 267, 17487-17490 (1992)). To measure CETP activity, [³H]CE-labeled HDL, LDL, CETP and assay buffer (50 mM tris(hydroxymethyl)aminomethane, pH 7.4; 150 mM sodium chloride; 2 mM ethylenediamine-tetraacetic acid; 1% bovine serum albumin) are incubated in a volume of 200 μ l, for 2 hours at 37°C in 96 well plates. LDL is differentially precipitated by the addition of 50 μ l of 1% (w/v) dextran sulfate/0.5 M magnesium chloride, mixed by vortex, and incubated at room temperature for 10 minutes. The solution (200 μ l) is transferred to a filter plate (Millipore). After filtration, the radioactivity present in the precipitated LDL is measured by liquid scintillation counting. Correction for non-specific transfer or precipitation is made by including samples that do not contain CETP. The rate of [³H]CE transfer using this assay is linear with respect to time and CETP concentration, up to 25-30% of [³H]CE transferred.

The potency of test compounds can be determined by performing the above described assay in the presence of varying concentrations of the test compounds and determining the concentration required for 50% inhibition of transfer of [³H]CE from HDL to LDL. This value is defined as the IC₅₀. The IC₅₀ values determined from this assay are accurate when the IC₅₀ is greater than 10 nM. In the case where compounds have greater inhibitory potency,

accurate measurements of IC_{50} may be determined using longer incubation times (up to 18 hours) and lower final concentrations of CETP (< 50 nM).

5

Inhibition of CETP Activity In Vivo.

Inhibition of CETP activity by a test compound can be determined by administering the compound to an animal by intravenous injection or oral gavage, measuring the amount
10 of transfer of tritium-labeled cholesteryl ester ($[^3H]CE$) from HDL to VLDL and LDL particles, and comparing this amount of transfer with the amount of transfer observed in control animals.

Male golden Syrian hamsters are maintained on a diet
15 of chow containing 0.24% cholesterol for at least two weeks prior to the study. For animals receiving intravenous dosing, immediately before the experiment,
~~animals are anesthetized with pentobarbital. Anesthesia~~
is maintained throughout the experiment. In-dwelling
20 catheters are inserted into the jugular vein and carotid artery. At the start of the experiment all animals will receive 0.2 ml of a solution containing $[^3H]CE$ -HDL into the jugular vein. $[^3H]CE$ -HDL is a preparation of human HDL containing tritium-labeled cholesteryl ester, and is
25 prepared according to the method of Glenn et al. (Meth. Enzymol., 263, 339-351 (1996)). Test compound is dissolved as a 80 mM stock solution in vehicle (2% ethanol: 98% PEG 400, Sigma Chemical Company, St. Louis, Missouri, USA) and administered either by bolus injection
30 or by continuous infusion. Two minutes after the $[^3H]CE$ -HDL dose is administered, animals receive 0.1 ml of the test solution injected into the jugular vein. Control animals receive 0.1 ml of the intravenous vehicle solution

without test compound. After 5 minutes, the first blood samples (0.5 ml) are taken from the carotid artery and collected in standard microtainer tubes containing ethylenediamine tetraacetic acid. Saline (0.5 ml) is
5 injected to flush the catheter and replace blood volume. Subsequent blood samples are taken at two hours and four hours by the same method. Blood samples are mixed well and kept on ice until the completion of the experiment. Plasma is obtained by centrifugation of the blood samples
10 at 4° C. The plasma (50 μ l) is treated with 5 μ l of precipitating reagent (dextran sulfate, 10 g/l; 0.5 M magnesium chloride) to remove VLDL/LDL. After centrifugation, the resulting supernatant (25 μ l) containing the HDL is analyzed for radioactivity using a
15 liquid scintillation counter.

The percentage [3 H]CE transferred from HDL to LDL and VLDL (% transfer) is calculated based on the total
~~radioactivity in equivalent plasma samples before~~
precipitation. Typically, the amount of transfer from HDL
20 to LDL and VLDL in control animals is 20% to 35% after 4 hours.

Alternatively, conscious, non-anesthetized animals can receive an oral gavage dose of test compound as a suspension in 0.1% methyl cellulose in water. At a time
25 determined for each compound at which plasma levels of the test substance reach their peak (C_{max}) after oral dosing, the animals are anesthetized with pentobarbital and then dosed with 0.2 ml of a solution containing [3 H]CE-HDL into the jugular vein as described above. Control animals
30 receive 0.25 ml of the vehicle solution without test compound by oral gavage. After 4 hours, the animals are sacrificed, blood samples are collected, and the

percentage [^3H]CE transferred from HDL to LDL and VLDL (% transfer) is assayed as described above.

Alternatively, inhibition of CETP activity by a test compound can be determined by administering the compound to mice that have been selected for expression of human CETP (hCETP) by transgenic manipulation (hCETP mice). Test compounds can be administered by intravenous injection, or oral gavage and the amount of transfer of tritium-labeled cholesteryl ester ([^3H]CE) from HDL to VLDL and LDL particles is determined, and compared to the amount of transfer observed in control animals. C57Bl/6 mice that are homozygous for the hCETP gene are maintained on a high fat chow diet, such as TD 88051, as described by Nishina et al. (J Lipid Res., 31, 859-869 (1990)) for at least two weeks prior to the study. Mice receive an oral gavage dose of test compound as a suspension in 0.1% methyl cellulose in water or an intravenous bolus ~~injection of test compound in 10% ethanol and 90%~~ polyethylene glycol. Control animals receive the vehicle solution without test compound by oral gavage or by an intravenous bolus injection. At the start of the experiment all animals receive 0.05 ml of a solution containing [^3H]CE-HDL into the tail vein. [^3H]CE-HDL is a preparation of human HDL containing tritium-labeled cholesteryl ester, and is prepared according to the method of Glenn et al. (Meth. Enzymol., 263, 339-351 (1996)). After 30 minutes, the animals are exsanguinated and blood collected in standard microtainer tubes containing ethylenediamine tetraacetic acid. Blood samples are mixed well and kept on ice until the completion of the experiment. Plasma is obtained by centrifugation of the blood samples at 4° C. The plasma is separated and analyzed by gel filtration chromatography and the relative

proportion of [^3H]CE in the VLDL, LDL and HDL regions is determined.

The percentage [^3H]CE transferred from HDL to LDL and VLDL (% transfer) is calculated based on the total
5 radioactivity in equivalent plasma samples before precipitation. Typically, the amount of transfer from HDL to LDL and VLDL in control animals is 20% to 35% after 30 min.

10 Intestinal Cholesterol Absorption Assay

A variety of compounds are shown to inhibit cholesterol absorption from the intestinal tract. These compounds lower serum cholesterol levels by reducing intestinal absorption of cholesterol from both exogenous
15 sources (dietary cholesterol) and endogenous cholesterol (secreted by the gall bladder into the intestinal tract).

In hamsters the use of a dual-isotope plasma ratio
~~method to measure intestinal cholesterol absorption has~~
been refined and evaluated as described by Turley et al.
20 (J. Lipid Res. 35, 329-339 (1994), herein incorporated by reference).

Male hamsters weighing 80-100 g are given food and water ad libitum in a room with 12 hour alternating periods of light and dark. Four hours into the light
25 period, each hamster is administered first an intravenous dose of 2.5 μCi of [$1,2\text{-}^3\text{H}$]cholesterol suspended in Intralipid (20%) and then an oral dose of [$4\text{-}^{14}\text{C}$]cholesterol in an oil of medium chain triglycerides (MCT). The i.v. dose is given by injecting a 0.4 ml volume
30 of the Intralipid mixture into the distal femoral vein. The oral dose is given by gavaging a 0.6 ml volume of the MCT oil mixture introduced intragastrically via a polyethylene tube. After 72 hours the hamsters are bled

and the amount of ^3H and ^{14}C in the plasma and in the original amount of label administered are determined by liquid scintillation spectrometry. The cholesterol absorption will be calculated based on the following
5 equation:

Percent cholesterol absorbed =

$$\frac{\% \text{ of oral dose per ml of 72 hour plasma sample}}{\% \text{ of i.v. dose per ml of 72 hour plasma sample}} \times 100$$

10

Microsomal triglyceride transfer protein (MTP) assay:

MTP can be purified from liver tissue or cultured
15 cells (e.g. HepG2 cells) using standard methods as described by Ohringer et al. (Acta Crystallogr. D52, 224-225 (1996), herein incorporated by reference).

~~Subsequent analysis of MTP activity can be performed~~
as described by Jamil et al. (Proc. Natl. Acad. Sci. 93,
20 11991-11995 (1996), herein incorporated by reference).

The basis of this assay is to measure the transfer of labeled triglycerides from a population of donor vesicles to a population of acceptor vesicles in the presence of MTP. Inhibitors of MTP can be evaluated by adding them to
25 the mixture prior to the introduction of MTP. Donor vesicles are to be prepared by sonication of an aqueous mixture of egg phospholipids, cardiolipin, ^3H -labeled phospholipid and ^{14}C -labeled triglycerides. Acceptor
30 vesicles are to be prepared by sonication of an aqueous mixture of egg phospholipids. The vesicle solutions are mixed together, with or without added MTP inhibitors, and MTP is to be added to initiate the transfer reaction. The assay will be terminated after 60 minutes by addition of

0.5 ml of DE-52 cellulose followed by centrifugation to pellet the donor molecules. The amount of ^3H and ^{14}C in the pellet and in the original amount of label in the mixture will be determined by liquid scintillation spectrometry. The lipid transfer rate will be calculated based on first order kinetics using the expression:

$$[S] = [S]_0 e^{-kt}$$

where $[S]_0$ and $[S]$ are the fractions of ^{14}C label in the donor membrane pellet at times 0 and t, respectively, and the term k is the fraction of label transferred per unit time.

15 Plasma Lipids Assay in Rabbits

Plasma lipids can be assayed using standard methods as reported by J.R. Schuh et al., J. Clin. Invest., 91, 1453-1458 (1993), herein incorporated by reference. Groups of male, New Zealand white rabbits are placed on a standard diet (100g/day) supplemented with 0.3% cholesterol and 2% corn oil (Zeigler Brothers, Inc., Gardners, PA). Water is available ad lib. Groups of control and treated animals are killed after 1 and 3 months of treatment. Tissues are removed for characterization of atherosclerotic lesions. Blood samples are to be taken for determination of plasma lipid concentrations.

Plasma Lipids

Plasma for lipid analysis is to be obtained by withdrawing blood from the ear vein into EDTA-containing tubes (Vacutainer; Becton Dickenson & Co., Rutherford,

NJ), followed by centrifugal separation of the cells. Total cholesterol was determined enzymatically, using the cholesterol oxidase reaction (C.A. Allain et al., Clin. Chem., 20, 470-475 (1974), herein incorporated by
5 reference). HDL cholesterol was also measured enzymatically, after selective precipitation of LDL and VLDL by dextran sulfate with magnesium (G.R. Warnick et al., Clin. Chem., 28, 1379-1388 (1982), herein
incorporated by reference). Plasma triglyceride levels
10 will be determined by measuring the amount of glycerol released by lipoprotein lipase through an enzyme-linked assay (G. Bucolo et al., Clin. Chem., 19, 476-482 (1973), herein incorporated by reference).

15 **Atherosclerosis**

Animals are to be killed by pentobarbital injection. Thoracic aortas are rapidly removed, immersion fixed in 10% neutral buffered formalin, and stained with oil red O

(0.3%). After a single longitudinal incision along the
20 wall opposite the arterial ostia, the vessels are pinned open for evaluation of the plaque area. The percent plaque coverage is determined from the values for the total area examined and the stained area, by threshold analysis using a true color image analyzer (Videometric
25 150; American Innovision, Inc., San Diego, CA) interfaced to a color camera (Toshiba 3CCD) mounted on a dissecting microscope. Tissue cholesterol will be measured enzymatically as described, after extraction with a chloroform/methanol mixture (2:1) according to the method
30 of Folch et al. (J. Biol. Chem., 226, 497-509 (1957), herein incorporated by reference).

In Vitro Vascular Response

The abdominal aortas are rapidly excised, after injection of sodium pentobarbital, and placed in oxygenated Krebs-bicarbonate buffer. After removal of perivascular tissue, 3-mm ring segments are cut, placed in
5 a 37°C muscle bath containing Krebs-bicarbonate solution, and suspended between two stainless steel wires, one of which is attached to a force transducer (Grass Instrument Co., Quincy, MA). Force changes in response to angiotensin II added to the bath will be recorded on a
10 chart recorder.

The examples herein can be performed by substituting the generically or specifically described therapeutic
15 compounds or inert ingredients for those used in the preceding examples.

The invention being thus described, it is apparent that the same can be varied in many ways. Such variations

are not to be regarded as a departure from the spirit and
20 scope of the present invention, and all such modifications and equivalents as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.

CLAIMS

What is claimed is:

1. A therapeutic combination comprising a first amount
5 of a nicotinic acid derivative compound and a
second amount of a cholesteryl ester transfer
protein inhibiting compound wherein the first
amount and the second amount together comprise an
anti-hyperlipidemic condition effective amount, an
10 anti-atherosclerotic condition effective amount, or
an anti-hypercholesterolemic condition effective
amount of the compounds.
2. The therapeutic combination of claim 1 wherein the
15 nicotinic acid derivative compound comprises
nicotinic acid.
3. The therapeutic combination of claim 1 wherein the
20 nicotinic acid derivative compound comprises
niceritrol.
4. The therapeutic combination of claim 1 wherein the
25 nicotinic acid derivative compound comprises
acipimox.
5. The therapeutic combination of claim 1 wherein the
combination comprises a composition comprising the
nicotinic acid derivative compound and the
cholesteryl ester transfer protein inhibiting
30 compound.
6. A method for the prophylaxis or treatment of a
hyperlipidemic condition comprising administering

to a patient in need thereof a combination in unit dosage form wherein the combination comprises a first amount of an nicotinic acid derivative compound and a second amount of a cholesteryl ester transfer protein inhibiting compound wherein the
5 first amount and the second amount together comprise an anti-hyperlipidemic condition effective amount of the compounds.

10 7. A method for the prophylaxis or treatment of an atherosclerotic condition comprising administering to a patient in need thereof a combination in unit dosage form wherein the combination comprises a
15 first amount of an nicotinic acid derivative compound and a second amount of a cholesteryl ester transfer protein inhibiting compound wherein the first amount and the second amount together comprise an anti-atherosclerotic condition
effective amount of the compounds.

20 8. A method for the prophylaxis or treatment of hypercholesterolemia comprising administering to a patient in need thereof a combination in unit dosage form wherein the combination comprises a
25 first amount of an nicotinic acid derivative compound and a second amount of a cholesteryl ester transfer protein inhibiting compound wherein the first amount and the second amount together comprise an anti-hypercholesterolemic condition
30 effective amount of the compounds.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/27942

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K45/06 A61P9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| A | WO 94 09774 A (MERCK) 11 May 1994 (1994-05-11) claims 1,6,9 page 1, line 6,7 ---- | 1,2,7,8 |
| A | DATABASE WPI Derwent Publications Ltd., London, GB; AN 1992-281671 XP002137557 BANYU PHARM: "Treatment agent for hypercholesterolaemia, hyperlipaemia or arteriosclerosis- contains squalene epoxidase inhibitor and other anti-hyperlipaemic agent." & JP 04 193836 A (BANYU PHARM), 13 July 1992 (1992-07-13) abstract ----- -/- | 1-3,6-8 |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 May 2000

Date of mailing of the international search report

23/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Peeters, J

PCT/US 99/27942

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Jonal Application No

PCT/US 99/27942

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9409774 A | 11-05-1994 | US 5256689 A | 26-10-1993 |
| | | AU 5538794 A | 24-05-1994 |
| JP 4193836 A | 13-07-1992 | NONE | |
| EP 818197 A | 14-01-1998 | DE 19627431 A | 15-01-1998 |
| | | AU 715101 B | 13-01-2000 |
| | | BG 101748 A | 30-04-1998 |
| | | BR 9703890 A | 03-11-1998 |
| | | CA 2209825 A | 08-01-1998 |
| | | CN 1174196 A | 25-02-1998 |
| | | CZ 9702144 A | 14-01-1998 |
| | | HR 970333 A | 30-04-1998 |
| | | HU 9701157 A | 30-03-1998 |
| | | JP 10167967 A | 23-06-1998 |
| | | NO 973143 A | 09-01-1998 |
| | | PL 320953 A | 19-01-1998 |
| | | SG 46781 A | 20-02-1998 |
| | | SK 92597 A | 06-05-1998 |
| | | US 5932587 A | 03-08-1999 |

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.